



Purchase Order No	Date	Fitter's Phone No.
Ordered by	Patient	Age
Street	Ship to	
City State Zip	Street	
Fitter	City State Zip	



STOCKINGS (Letters, Dotted Lines)


LEFT			RIGHT		
Girth	Length		Girth	Length	
I	F to I		I	F to I	
H	F to H		H	F to H	
G	F to G		G	F to G	
F	Floor to F		F	Floor to F	
E	Floor to E		E	Floor to E	
D	Floor to D		D	Floor to D	
X	Floor to X		X	Floor to X	
C	Floor to C		C	Floor to C	
O	Flex to heel tip	X	O	Flex to heel tip	X
B	Heel tip to B		B	Heel tip to B	
A	Heel tip to A		A	Heel tip to A	
BT	Heel tip to BT		BT	Heel tip to BT	



GARTER BELTS AND CHAPS (Numbers, Solid Lines)

1		<input type="checkbox"/> Left Leg
2		<input type="checkbox"/> Right Leg

Form F-030 IMF - for Use Without Paper Tapes



Please refer to our current price list for a description of these items

HOTLINE SERVICE

Guaranteed delivery in 4-5 business days, or less) **Additional Cost**

Sex: M F

REQUIRED Allergies:

Allergic to silicone? Yes, No, NA | Allergic to metal? Yes, No, NA

Knee Length: L R | **Thigh Length:** L R

Chaps: 300 L R | 307 | Both

WAIST LENGTH SUPPORTS:

See Form F-029

OPTIONS & EXTRAS:

● **Zipper:** Inside (or) Outside
 Open (or) Closed

● **Garment Color No:** _____

● **Toe Caps:** None/Open (or) Soft* (or) Self*
(*Indicate Foot Length in Diagram)

● **Reinforcement:** Heel | Knee Front
 Knee Back | Inner Thigh

● **Flex/Contracture Seam:** Instep | Knee Back

● **Elastic Band:** Regular 1" 2"
(or) Silicone 1" 2"

● **Other Option(s):** _____

INDICATIONS:

- 20-30 mm Hg:** Varicose veins (mild), arterial insufficiency with venous insufficiency.
- 22-28 mm Hg:** Burns — prevention of hypertrophic scars.
- 30-40 mm Hg:** Varicose veins (moderate), assist fluid return, leg fatigue, stasis dermatitis, postphlebotic syndrome, post surgical stripping of sclerosing, postfracture edema, prophylactic treatment of edema and phlebitis, lymphedema (moderate).
- 40-50 mm Hg:** Chronic venous insufficiency, stasis dermatitis (severe), lymphedema (severe), chronic venous insufficiency (severe), orthostatic hypotension (moderate).
- 50-60 mm Hg:** Orthostatic hypotension (severe), postthrombosis (severe), intractable edema.

Contraindications: Non-ambulatory use, severe arterial insufficiency, cutaneous infection, acute hypodermatitis, wet dermatitis, dermatitis.

Check this box if there are instructions on other side