


Purchase Order No	Date	Fitter's Phone No.
Ordered by	Patient	Age
Street	Ship to	
City	State	Zip
Fitter	City	State
		Zip



Please refer to our current price list for a description of these items


IMPORTANT:

Lengths to be taken from the back and from the floor barefooted to the girth points. Please measure carefully and complete all appropriate boxes.

NOTE:

One leg waist length supports with closed crotch can only be made in panty style, (Cat. No. 303). Measure girth of unsupported leg at fold of buttocks, and girth 5" below fold.

Pregnancy garments available only with closed crotch, recommended total length measured to F.



GIRTHS		LENGTHS	
F		Floor to F	
E		Floor to E	
D		Floor to D	
C		Floor to C	
L		Inseam	
R			

Height		Weight	
--------	--	--------	--

HOTLINE SERVICE

(Guaranteed delivery in 4-5 business days, or less) **Additional Cost**

Sex: M F

REQUIRED Allergies:

Allergic to silicone? Yes, No, NA | Allergic to metal? Yes, No, NA

WAIST LENGTH SUPPORTS ONLY:

One Leg (or) Bilateral

Crotch: Open (or) Closed (or) Fly Front

Pregnancy, month _____

OPTIONS & EXTRAS:

● **Zipper:**

Left Side (or) Left Off Center
 Right Side (or) Right Off Center

● **Garment Color No:** _____

● **Abdominal Panel:**

Double (or) Uplift (or)
 Reduced Pressure

● **Crotch:** Reinforced

● **Inner Thigh:** Reinforced

● **Elastic Band:** Regular 1" 2" (or)
Silicone 1" 2"

INDICATIONS:

- 20-30 mm Hg:** Varicose veins (mild), arterial insufficiency with venous insufficiency.
- 22-28 mm Hg:** Burns — prevention of hypertrophic scars.
- 30-40 mm Hg:** Varicose veins (moderate), assist fluid return, leg fatigue, stasis dermatitis, postphlebotic syndrome, post surgical stripping of sclerosing, postfracture edema, prophylactic treatment of edema and phlebitis, lymphedema (moderate).
- 40-50 mm Hg:** Chronic venous insufficiency, stasis dermatitis (severe), lymphedema (severe), chronic venous insufficiency (severe), orthostatic hypotension (moderate).
- 50-60 mm Hg:** Orthostatic hypotension (severe), postthrombosis (severe), intractable edema.

Contraindications: Non-ambulatory use, severe arterial insufficiency, cutaneous infection, acute hypodermatitis, wet dermatitis.